

# Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/EP04/013687

International filing date: 02 December 2004 (02.12.2004)

Document type: Certified copy of priority document

Document details: Country/Office: US  
Number: 60/526,609  
Filing date: 04 December 2003 (04.12.2003)

Date of receipt at the International Bureau: 17 March 2005 (17.03.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland  
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

PCT/EP200 4 / 0 1 3 6 8 7

01.03.05

PA 1280482

# THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office


February 08, 2005

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.

APPLICATION NUMBER: 60/526,609

FILING DATE: December 04, 2003

By Authority of the  
COMMISSIONER OF PATENTS AND TRADEMARKS



P. R. GRANT  
Certifying Officer

Mail Stop Patent Applica

PROVISIONAL APPLICATION COVER SHEET

PF55118/Za

This is a request for filing a PROVISIONAL APPLICATION under 35 USC 111(b).

1. INVENTOR(s)		
Last Name	First Name	Residence
OLOUMI-SADEGHI	Hassan	12105 Pawley's Mill Circle Raleigh, NC 27614 USA Citizen of USA
KUHN	David G.	1208 Dalgarn Drive Apex, NC 27502 USA Citizen of USA
VON DEYN	Wolfgang	An der Bleiche 24 67435 Neustadt GERMANY Citizen of Germany
ARMES	Nigel	8001 Kukui Court Raleigh, NC 27613 USA Citizen of Great Britain


22581 U.S. PTO  
60/526609



120403

2. TITLE OF THE INVENTION: **USE OF N-ARYLHYDRAZINE DERIVATIVES FOR COMBATING PESTS**
3. CORRESPONDENCE ADDRESS:  
Keil & Weinkauff  
1350 Connecticut Ave., N.W.  
Washington, D.C. 20036  
(202) 659-0100; (202) 659-0105-fax
4. ENCLOSED APPLICATION PARTS:  
☒ Specification 30 pages  
☐ Drawings — sheets  
☒ Claims 9 claims
5. METHOD OF PAYMENT  
☒ A check in the amount of \$160.00 is attached to cover the required Provisional filing fee.  
☒ The commissioner is hereby authorized to charge any deficiency in fees to Deposit Account 11.0345.
6. The invention was **not** made by an agency of the United States Government or under a contract with an agency of the United States Government.

Respectfully submitted,  
KEIL & WEINKAUF


  
Herbert B. Keil  
Reg. No. 18,967

HBK/kas

1350 Connecticut Ave., N.W.  
Washington, D.C. 20036  
(202)659-0100




The use of N-arylhydrazine derivatives for combating pests

The present invention relates to the use of hydrazine derivatives of formula I:



wherein

Q is



$X^1$  is chlorine, bromine, or fluorine;

$R^1, R^2$  are each independently hydrogen,  $C_1$ - $C_{10}$ -alkyl,  $C_3$ - $C_{10}$ -alkenyl,  $C_3$ - $C_{10}$ -alkynyl, or  $C_3$ - $C_{12}$ -cycloalkyl,  $C_1$ - $C_6$ -alkylamino, di( $C_1$ - $C_6$ -alkyl)-amino,  $C_1$ - $C_6$ -alkylcarbonylamino,  $C_1$ - $C_6$ -alkylsulfonyl, or  $C_1$ - $C_6$ -alkylsulfinyl, wherein the carbon atoms in these groups may be substituted with

1 to 3 halogen, hydroxy, nitro, cyano, amino, mercapto,  $C_1$ - $C_6$ -alkoxy,  $C_1$ - $C_6$ -haloalkoxy,  $C_1$ - $C_6$ -alkylthio,  $C_1$ - $C_6$ -haloalkylthio,  $C_1$ - $C_6$ -alkylsulfonyl,  $C_1$ - $C_6$ -alkylsulfinyl,  $C_1$ - $C_6$ -haloalkylsulfonyl,  $C_1$ - $C_6$ -haloalkylsulfinyl, or  $C_3$ - $C_6$ -cycloalkyl which may be substituted with 1 to 3  $R^{\#}$  groups, or


$R^{\#}$  is halogen, cyano, nitro, hydroxy, mercapto, amino,  $C_1$ - $C_6$ -alkoxy,  $C_2$ - $C_6$ -alkenyloxy,  $C_2$ - $C_6$ -alkynyloxy,  $C_1$ - $C_6$ -haloalkoxy,  $C_1$ - $C_6$ -alkylthio, or  $C_1$ - $C_6$ -haloalkylthio,  $C_1$ - $C_6$ -alkylsulfonyl,  $C_1$ - $C_6$ -alkylsulfinyl,  $C_1$ - $C_6$ -alkylamino, di( $C_1$ - $C_6$ -alkyl)-amino,  $C_1$ - $C_6$ -alkylcarbonyl,  $C_1$ - $C_6$ -alkoxycarbonyl, or di( $C_1$ - $C_6$ )-alkylaminocarbonyl;

formyl,  $C_1$ - $C_6$ -alkylcarbonyl,  $C(=O)NR^aR^b$ ,  $CO_2R^c$ ,  $R^d$ ,  $R^e$ , phenyl which may be substituted with 1 to 3  $R^{\#}$  groups, or pyridyl which may be substituted with 1 to 3  $R^{\#}$  groups,

$R^a, R^b, R^c$  are each independently hydrogen or  $C_1$ - $C_4$ -alkyl which may be substituted with 1 to 3 groups  $R^{\#}$ ;

$R^d$  is  $NR^1R^2$  or

2



$R^1, R^1$  are each independently hydrogen or  $C_1$ - $C_4$ -alkyl which may be substituted with 1 to 3 groups  $R^H$ ;

5


$p, m$  are each independently 0, 1, 2, or 3, with the proviso that  $p$  and  $m$  are not both 0.

$X$  is oxygen, sulfur, amino,  $C_1$ - $C_4$ -alkylamino, or phenylamino, or, if  $p$  is 0 then  $X$  can also be phenoxy or  $C_1$ - $C_6$ -alkoxy;

$r$  is 0 or 1;

10


$R^B$  is



15

$R^k, R^a$  are each independently hydrogen or  $C_1$ - $C_4$ -alkyl which may be substituted with 1 to 3 groups  $R^H$ ; or

$R^1$  and  $R^2$  may be taken together to form a ring represented by the structure



20

$p, m$  are 1, 2 or 3;

$X'$  is oxygen, sulfur, amino,  $C_1$ - $C_4$ -alkylamino, phenylamino, or methylene;

$Z$  is  $C_1$ - $C_4$ -alkyl or phenyl;

25

$R^3$  is hydrogen,  $C_1$ - $C_{10}$ -alkyl,  $C_2$ - $C_{10}$ -alkenyl,  $C_2$ - $C_{10}$ -alkynyl,  $C_3$ - $C_{12}$ -cycloalkyl, wherein the carbon atoms in these groups may be partially or fully halogenated or substituted with

30

1 to 3 cyano, nitro, hydroxy, mercapto, amino,  $C_1$ - $C_6$ -alkyl,  $C_3$ - $C_6$ -cycloalkyl,  $C_1$ - $C_6$ -alkoxy,  $C_1$ - $C_6$ -alkylamino, di( $C_1$ - $C_6$ -alkyl)-amino,  $C_1$ - $C_6$ -alkylthio,  $C_1$ - $C_6$ -alkylsulfonyl, or  $C_1$ - $C_6$ -alkylsulfinyl groups, wherein the carbon atoms in these groups may be substituted by

35

1 to 3 halogen atoms, a 5- to 6-membered aromatic ring system which may contain 1 to 4 heteroatoms selected from oxygen, sulfur and nitrogen and which may be substituted with any combination of 1 to 5 halogen atoms, 1 to 3  $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -alkylthio,  $C_1$ - $C_6$ -alkylsulfonyl,  $C_1$ - $C_6$ -

## 3

alkylsulfinyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, nitro, or cyano groups, wherein the carbon atoms in these groups may be substituted by 1 to 3 halogen atoms, or

5

phenoxy, which may be substituted with any combination of 1 to 5 halogen atoms, 1 to 3 C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-alkylthio, C<sub>1</sub>-C<sub>6</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, nitro, or cyano groups, wherein the carbon atoms in these groups may be substituted by 1 to 3 halogen atoms, or

10

a 3- to 6-membered saturated or partially unsaturated ring system which contains 1 to 3 heteroatoms selected from oxygen, sulfur and nitrogen and which may be substituted with any combination of 1 to 5 halogen atoms, 1 to 3 C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-alkylthio, C<sub>1</sub>-C<sub>6</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, nitro, or cyano groups, wherein the carbon atoms in these groups may be substituted by 1 to 3 halogen atoms,

15

a 3- to 6-membered saturated or partially unsaturated ring system which contains 1 to 3 heteroatoms selected from oxygen, sulfur and nitrogen and which is unsubstituted or substituted with any combination of 1 to 5 halogen atoms, 1 to 3 C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-alkylthio, C<sub>1</sub>-C<sub>6</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, C<sub>1</sub>-C<sub>6</sub>-haloalkoxy, nitro, or cyano groups, wherein the carbon atoms in these groups may be substituted by 1 to 3 halogen atoms;

20

25 R, R<sup>4</sup> are each independently hydrogen or C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkylaminocarbonyl, or di(C<sub>1</sub>-C<sub>6</sub>-alkyl)-aminocarbonyl, wherein the carbon atoms in the these groups may be substituted with 1 to 3 groups R<sup>#</sup>;

A is C-R<sup>5</sup> or N;

B is C-R<sup>6</sup> or N;

30 W is C-R<sup>7</sup> or N;

with the proviso that one of A, B and W is other than N;

35

R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> are each independently hydrogen, halogen, nitro, cyano, amino, mercapto, hydroxy, C<sub>1</sub>-C<sub>10</sub>-alkyl, C<sub>2</sub>-C<sub>10</sub>-alkenyl, C<sub>2</sub>-C<sub>10</sub>-alkynyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, C<sub>1</sub>-C<sub>6</sub>-alkylamino, di(C<sub>1</sub>-C<sub>6</sub>-alkyl)-amino, C<sub>1</sub>-C<sub>6</sub>-alkylthio, C<sub>1</sub>-C<sub>6</sub>-alkylsulfonyl, or C<sub>1</sub>-C<sub>6</sub>-alkylsulfinyl, wherein the carbon atoms in these groups may be substituted with 1 to 3 groups R<sup>#</sup>

40

a 5- to 6-membered aromatic ringsystem which may contain 1 to 4 heteroatoms selected from oxygen, sulfur and nitrogen and which may be substituted with any combination of 1 to 5 halogen atoms, 1 to 3 C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>1</sub>-C<sub>6</sub>-alkylthio, C<sub>1</sub>-C<sub>6</sub>-haloalkylthio, C<sub>1</sub>-C<sub>6</sub>-

## 4

alkylsulfonyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>6</sub>-haloalkylsulfonyl, C<sub>1</sub>-C<sub>6</sub>-haloalkylsulfinyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, C<sub>1</sub>-C<sub>6</sub>-haloalkoxy, mercapto, hydroxy, amino, nitro, or cyano groups, wherein the carbon atoms in these groups may be substituted with 1 to 3 groups R<sup>#</sup>;

5 Y is hydrogen, halogen, cyano, nitro, amino, hydroxy, mercapto, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>2</sub>-C<sub>10</sub>-alkenyl, C<sub>2</sub>-C<sub>10</sub>-alkynyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, C<sub>1</sub>-C<sub>6</sub>-alkylamino, di(C<sub>1</sub>-C<sub>6</sub>)-alkylamino, C<sub>1</sub>-C<sub>6</sub>-alkylthio, C<sub>1</sub>-C<sub>6</sub>-alkylsulfonyl, or C<sub>1</sub>-C<sub>6</sub>-alkylsulfinyl, wherein the carbon atoms in these groups may be substituted with 1 to 3 groups R<sup>#</sup>;

10

n is 0, 1, or 2;

or the enantiomers or diastereomers, salts or esters thereof,

15 for combating non-crop pests.

The invention also relates to a method for controlling pests selected from the above orders comprising contacting the pests or their food supply, habitat or breeding grounds with a pesticidally effective amount of compounds of formula I.

20

The invention also relates to compositions for controlling pests of the above orders.

Typical problems arising with the use of presently available non-crop pest controllers with a broad spectrum of activity such as pyrethroids are e.g. resistance of pests or unfavorable toxicological properties. Accordingly, there is a need to provide new and improved non-crop pest controllers that overcome these problems.

25

It is therefore an object of the present invention to provide new non-crop pest controllers exhibiting an enhanced pesticidal spectrum of action.

30

It is also an object of the present invention to provide a method for controlling pests selected from the above orders.

We have found that these objects are achieved by use of compounds of formula I and compositions comprising them.

35

The hydrazine derivatives of formula I which can be used according to the invention are known from EP-A 604 798. This document relates to plant protection in the agricultural filed and discloses the insecticidal and acaricidal activity of compounds of formula I against crop pests of the Coleoptera, Lepidoptera and Acarina orders.

40

## 5

The compounds of formula I can be prepared according to preparation methods described or referenced in EP-A 604 798 or modifications thereof.

5 In the definition of formula I shown above, the substituents have the following meanings:

"Halogen" will be taken to mean fluoro, chloro, bromo and iodo.

10 The term "alkyl" as used herein refers to a branched or unbranched saturated hydrocarbon group having 1 to 10 carbon atoms, especially C<sub>1</sub>-C<sub>6</sub>-alkyl such as methyl, ethyl, propyl, 1-methylethyl, butyl, 1-methylpropyl, 2-methylpropyl, 1,1-dimethylethyl, pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 2,2-dimethylpropyl, 1-ethylpropyl, hexyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 15 2,2-dimethylbutyl, 2,3-dimethylbutyl, 3,3-dimethylbutyl, 1-ethylbutyl, 2-ethylbutyl, 1,1,2-trimethylpropyl, 1,2,2-trimethylpropyl, 1-ethyl-1-methylpropyl and 1-ethyl-2-methylpropyl.

20 The term "haloalkyl" as used herein refers to a straight-chain or branched alkyl groups having 1 to 10 carbon atoms (as mentioned above), where some or all of the hydrogen atoms in these groups may be replaced by halogen atoms as mentioned above, for example C<sub>1</sub>-C<sub>2</sub>-haloalkyl, such as chloromethyl, bromomethyl, dichloromethyl, tri-chloromethyl, fluoromethyl, difluoromethyl, trifluoromethyl, chlorofluoromethyl, dichloro-fluoromethyl, chlorodifluoromethyl, 1-chloroethyl, 1-bromoethyl, 1-fluoroethyl, 2-25 fluoroethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl, 2-chloro-2-fluoroethyl, 2-chloro-2,2-difluoroethyl, 2,2-dichloro-2-fluoroethyl, 2,2,2-trichloroethyl and pentafluoroethyl;

30 "Alkoxy" refers to straight-chain or branched alkyl group having 1 to 4 or 6 carbon atoms (as mentioned above) bonded through an oxygen linkage, at any bond in the alkyl group. Examples include methoxy, ethoxy, propoxy, and isopropoxy.

35 Likewise, the terms "alkylthio", "alkylamino", "dialkylamino", "alkylsulfonyl", and alkyl-sulfinyl" refer to straight-chain or branched alkyl group having 1 to 4 or 6 carbon atoms (as mentioned above) bonded through a sulfur-, -NH-, -N-, -S(=O)<sub>2</sub>-, or S(=O)- linkage, respectively.

40 The term "alkenyl" as used herein intends a branched or unbranched unsaturated hydrocarbon group having 3 to 10 carbon atoms and a double bond in any position, such as C<sub>3</sub>-C<sub>6</sub> alkenyl such as 1-propenyl, 2-propenyl, 1-methyl-ethenyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-methyl-1-propenyl, 2-methyl-1-propenyl, 1-methyl-2-propenyl, 2-methyl-2-propenyl; 1-pentenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 1-methyl-1-butenyl, 2-methyl-1-butenyl, 3-methyl-1-butenyl, 1-methyl-2-butenyl, 2-methyl-2-butenyl, 3-



## 6

methyl-2-butenyl, 1-methyl-3-butenyl, 2-methyl-3-butenyl, 3-methyl-3-butenyl, 1,1-dimethyl-2-propenyl, 1,2-dimethyl-1-propenyl, 1,2-dimethyl-2-propenyl, 1-ethyl-1-propenyl, 1-ethyl-2-propenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 5-hexenyl, 1-methyl-1-pentenyl, 2-methyl-1-pentenyl, 3-methyl-1-pentenyl, 4-methyl-1-pentenyl, 1-methyl-2-pentenyl, 2-methyl-2-pentenyl, 3-methyl-2-pentenyl, 4-methyl-2-pentenyl, 1-methyl-3-pentenyl, 2-methyl-3-pentenyl, 3-methyl-3-pentenyl, 4-methyl-3-pentenyl, 1-methyl-4-pentenyl, 2-methyl-4-pentenyl, 3-methyl-4-pentenyl, 4-methyl-4-pentenyl, 1,1-dimethyl-2-butenyl, 1,1-dimethyl-3-butenyl, 1,2-dimethyl-1-butenyl, 1,2-dimethyl-2-butenyl, 1,2-dimethyl-3-butenyl, 1,3-dimethyl-1-butenyl, 1,3-dimethyl-2-butenyl, 1,3-dimethyl-3-butenyl, 2,2-dimethyl-3-butenyl, 2,3-dimethyl-1-butenyl, 2,3-dimethyl-2-butenyl, 2,3-dimethyl-3-butenyl, 3,3-dimethyl-1-butenyl, 3,3-dimethyl-2-butenyl, 1-ethyl-1-butenyl, 1-ethyl-2-butenyl, 1-ethyl-3-butenyl, 2-ethyl-1-butenyl, 2-ethyl-2-butenyl, 2-ethyl-3-butenyl, 1,1,2-trimethyl-2-propenyl, 1-ethyl-1-methyl-2-propenyl, 1-ethyl-2-methyl-1-propenyl and 1-ethyl-2-methyl-2-propenyl;

15

"Cycloalkyl" refers to a monocyclic 3- to 6-, 8-, 10- or 12-membered saturated carbon atom rings, e.g. C<sub>3</sub>-C<sub>8</sub>-cycloalkyl such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl.

20

A 5- to 6-membered aromatic ring system containing 1 to 4 heteroatoms selected from oxygen, sulfur and nitrogen, intends e.g. 5-membered hetaryl, containing 1 to 4 nitrogen atoms or 1 to 3 nitrogen atoms and 1 sulfur or oxygen atom, e.g. furyl, thienyl, pyrrolyl, isoxazolyl, isothiazolyl, pyrazolyl, oxazolyl, thiazolyl, imidazolyl, oxadiazolyl, thiadiazolyl, oxadiazolyl, triazolyl, and tetrazolyl; or 6-membered hetaryl, containing 1 to 4 nitrogen atoms or 1 to 3 nitrogen atoms and 1 sulfur or oxygen atom, e.g. 2-pyridinyl, 3-pyridinyl, 4-pyridinyl, 3-pyridazinyl, 4-pyridazinyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 2-pyrazinyl, 1,3,5-triazin-2-yl and 1,2,4-triazin-3-yl;

25

a 3- to 6-membered saturated or partially unsaturated ring system which contains 1 to 3 heteroatoms selected from oxygen, sulfur and nitrogen intends e.g. a saturated 3- to 6-membered ringsystem containing 1 to 3 heteroatoms selected from nitrogen and oxygen, such as aziridine, pyrrolidine, tetrahydrofuran, tetrahydropyran, or piperidine.

30

With respect to the intended use of the compounds of formula I, particular preference is given to the following meanings of the substituents, in each case on their own or in combination:

35

Preference is given to compounds of formula I wherein A denotes C-R<sup>5</sup>.

40

Moreover, preference is given to compounds of formula I wherein B denotes C-R<sup>6</sup>.

Preference is also given to compounds of formula I wherein W denotes C-R<sup>7</sup>.

Particular preference is given to compounds wherein A denotes C-R<sup>5</sup>, B denotes C-R<sup>6</sup>, and W denotes C-R<sup>7</sup>.

- 5 Moreover, preference is given to compounds of formula I wherein R<sup>5</sup> is halogen or C<sub>1</sub>-C<sub>6</sub>-haloalkyl, with halogen, especially chlorine, being most preferred.

Preference is also given to compounds of formula I wherein R<sup>6</sup> is hydrogen or halogen, especially hydrogen.

10

Preference is further given to compounds of formula I wherein R<sup>7</sup> is halogen or C<sub>1</sub>-C<sub>6</sub>-haloalkyl, preferably C<sub>1</sub>-C<sub>6</sub>-haloalkyl, especially trifluoromethyl.

- 15 Moreover, preference is given to compounds of formula I wherein Y is halogen or C<sub>1</sub>-C<sub>6</sub>-haloalkyl. Particular preference is given to compounds of formula I wherein Y is halogen, especially chlorine.

Preference is given to compounds of formula I wherein n is 1.

- 20 Preference is also given to compounds of formula I wherein Q denotes -N=[C(NR<sup>1</sup>R<sup>2</sup>)R<sup>3</sup>].

Moreover, preference is given to compounds of formula I wherein X<sup>1</sup> is chlorine.

- 25 Preference is also given to compounds of formula I wherein R denotes C<sub>1</sub>-C<sub>6</sub>-alkyl or hydrogen.

Preference is also given to compounds of formula I wherein R<sup>1</sup> and R<sup>2</sup> each independently are hydrogen, C<sub>1</sub>-C<sub>10</sub>-alkyl, or C<sub>3</sub>-C<sub>10</sub>-cycloalkyl.

30

Moreover, preference is given to compounds of formula I wherein R<sup>1</sup> and R<sup>2</sup> each independently are hydrogen, C<sub>1</sub>-C<sub>4</sub>-alkyl, or C<sub>3</sub>-C<sub>6</sub>-cycloalkyl.

- 35 Moreover, preference is given to compounds of formula I wherein R<sup>3</sup> is unsubstituted C<sub>1</sub>-C<sub>10</sub>-alkyl or C<sub>3</sub>-C<sub>10</sub>-cycloalkyl, which may be substituted with 1 to 5 halogen atoms and/or 1-3 C<sub>1</sub>-C<sub>6</sub>-alkyl groups.


Particularly preferred are compounds of formula I wherein R<sup>3</sup> is tert.-butyl.

- 40 Moreover, particularly preferred are compounds of formula I wherein R<sup>3</sup> is cyclopropyl which may be substituted with C<sub>1</sub>-C<sub>6</sub>-alkyl or halogen, especially 1-methyl-2,2-dichlorocyclopropyl.


Moreover, preference is given to compounds of formula I wherein R<sup>4</sup> is hydrogen or C<sub>1</sub>-C<sub>6</sub>-alkyl.

- 5 Particular preference is given to N-ethyl-2,2-dimethylpropionamide,2-(2,6-dichloro- $\alpha$ ,  $\alpha$ ,  $\alpha$ -trifluoro-p-tolyl)hydrazone and N-Ethyl-2,2-dichloro-1-methylcyclopropanecarboxamide, 2-(2,6-dichloro- $\alpha$ , $\alpha$ , $\alpha$ -trifluoro-p-tolyl)hydrazone.

- 10 Furthermore, particular preference with respect to the use in the present invention is given to the compound of formula I-1 (N-ethyl-2,2-dimethylpropionamide,2-(2,6-dichloro- $\alpha$ ,  $\alpha$ ,  $\alpha$ -trifluoro-p-tolyl)-hydrazone):



- 15 Moreover, particular preference with respect to the use in the present invention is given to the compound of formula I-2 (N-Ethyl-2,2-dichloro-1-methylcyclopropanecarboxamide, 2-(2,6-dichloro- $\alpha$ , $\alpha$ , $\alpha$ -tri-fluoro-p-tolyl)hydrazone):



- 20 The compounds of formula I are suitable for combating non-crop pests in the protection of stored products and of materials, in the hygiene field, and also in forestry and agriculture. They are active against all or some stages of development.

- Non-crop pests are commonly understood as pests from the classes Chilopoda and Diplopoda, from the orders Isoptera, Blattaria (Blattodea), Diptera, Dermaptera, Hemiptera, Thysanoptera, Hymenoptera, Orthoptera, Siphonaptera, Thysanura, Phthiraptera, Mesostigmata, Prostigmata, Astigmata, and Ixodida and from the families Saturniidae,  
25 Pyralidae, Gelechiidae, Meloidae.

The compounds of the formula I are especially suitable for efficiently combating the following pests:

- 30 centipedes (*Chilipoda*), e.g. *Scutigera coleoptrata*,  
millipedes (*Diplopoda*), e.g. *Narceus spp.*,

## 9

termites (Isoptera), e.g. *Calotermes flavicollis*, *Leucotermes flavipes*, *Heterotermes aureus*, *Reticulitermes flavipes*, *Reticulitermes virginicus*, *Reticulitermes lucifugus* and *Termes natalensis*, *Coptotermes formosanus*

- 5 cockroaches (Blattaria - Blattodea), e.g. *Blattella germanica*, *B. asahinae*, *Periplaneta americana*, *Periplaneta japonica*, *Periplaneta brunnea*, *Periplaneta fuliginosa*, *P. australasiae*, and *Blatta orientalis*

- flies, mosquitoes (Diptera), e.g. *Aedes aegypti*, *Aedes albopictus*, *Aedes vexans*, *Anastrepha ludens*, *Anopheles maculipennis*, *Anopheles crucians*, *An. albimanus*, *An. Gambiae*, *An. freeborni*, *An. leucosphyrus*, *An. minimus*, *An. quadrimaculatus*, *Calliphora vicina*, *Ceratitis capitata*, *Chrysomya bezziana*, *Chrysomya hominivorax*, *Chrysomya macellaria*, *Chrysomya bezziana*, *Chrysops discalis*, *C. silacea*, *C. atlanticus*, *Cochliomyia hominivorax*, *Contarinia sorghicola*, *Cordylobia anthropophaga*, *Culicoides furens*, *Culex pipiens*, *Culex nigripalpus*, *C. quinquefasciatus*, *C. tarsalis*, *Culiseta inornata*, *C. melanura*, *Dacus cucurbitae*, *Dacus oleae*, *Dasineura brassicae*, *Dermatobia hominis*, *Fannia canicularis*, *Gasterophilus intestinalis*, *Glossina morsitans*, *Glossina palpalis*, *G. fuscipes*, *G. tachinoides*, *Haematobia irritans*, *Haplodiplosis equestris*, *Hippelates spp.*, *Hylemyia platura*, *Hypoderma lineata*, *Leptoconops torrens*, *Liriomyza sativae*, *Liriomyza trifolii*, *Lucilia caprina*, *Lucilia cuprina*, *Lucilia sericata*, *Lycoria pectoralis*, *Mansonella titillans*, *Mayetiola destructor*, *Musca domestica*, *Muscina stabulans*, *Oestrus ovis*, *Oscinella frit*, *Pegomya hysocyami*, *Phorbia antiqua*, *Phorbia brassicae*, *Phorbia coarctata*, *Phlebotomus argentipes*, *Psorophora columbiae*, *P. discolor*, *Prosimulium mixtum*, *Rhagoletis cerasi*, *Rhagoletis pomonella*, *Sarcophaga haemorrhoidalis*, *Sarcophaga sp.*, *Simulium vittatum*, *Stomoxys calcitrans*, *Tabanus bovinus*, *Tabanus atratus*, *T. lineola*, *T. similis*, *Tipula oleracea*, and *Tipula paludosa*
- 25

Earwigs (Dermaptera), e.g. *forficula auricularia*,

- 30 true bugs (Hemiptera), e.g. *Acrosternum hilare*, *Blissus leucopterus*, *Cyrtopeltis notatus*, *Dysdercus cingulatus*, *Dysdercus intermedius*, *Eurygaster integriceps*, *Euschistus impictiventris*, *Leptoglossus phyllopus*, *Lygus lineolaris*, *Lygus pratensis*, *Nezara viridula*, *Piesma quadrata*, *Solubea insularis*, *Thyanta perditor*, *Acyrtosiphon onobrychis*, *Adelges laricis*, *Aphidula nasturtii*, *Aphis fabae*, *Aphis forbesi*, *Aphis pomi*, *Aphis gossypii*, *Aphis grossulariae*, *Aphis schneideri*, *Aphis spiraecola*, *Aphis sambuci*, *Acyrtosiphon pisum*, *Aulacorthum solani*, *Brachycaudus cardui*, *Brachycaudus helichrysi*, *Brachycaudus persicae*, *Brachycaudus prunicola*, *Brevicoryne brassicae*, *Capitophorus horni*, *Cerosiphia gossypii*, *Chaetosiphon fragaefolii*, *Cryptomyzus ribis*, *Dreyfusia nordmanniana*, *Dreyfusia piceae*, *Dysaphis radicola*, *Dysaulacorthum pseudosolani*, *Dysaphis plantaginea*, *Dysaphis pyri*, *Empoasca fabae*, *Hyalopterus pruni*, *Hyperomyzus lactucae*, *Macrosiphum avenae*, *Macrosiphum euphorbiae*, *Macrosiphon rosae*, *Megoura viciae*, *Melanaphis pyramis*, *Metopolophium dirhodum*, *Myzodes persicae*,
- 40

## 10

- Myzus ascalonicus*, *Myzus cerasi*, *Myzus varians*, *Nasonovia ribis-nigri*, *Nilaparvata lugens*, *Pemphigus bursarius*, *Perkinsiella saccharicida*, *Phorodon humuli*, *Psylla mali*, *Psylla piri*, *Rhopalomyzus ascalonicus*, *Rhopalosiphum maidis*, *Rhopalosiphum padi*, *Rhopalosiphum insertum*, *Sappaphis mala*, *Sappaphis mali*, *Schizaphis graminum*,  
5 *Schizoneura lanuginosa*, *Sitobion avenae*, *Trialeurodes vaporariorum*, *Toxoptera aurantiiand*, *Viteus vitifolii*, *Cimex lectularius*, *C. hemipterus*, *Reduvius senilis*, *Triatoma* spp., and *Arilus critatus*.
- thrips (*Thysanoptera*), e.g. *Frankliniella fusca*, *Frankliniella occidentalis*, *Frankliniella tritici*, *Scirtothrips citri*, *Thrips oryzae*, *Thrips palmi* and *Thrips tabaci*,  
10
- ants, bees, wasps, sawflies (*Hymenoptera*), e.g. *Athalia rosae*, *Atta cephalotes*, *Atta sexdens*, *Atta texana*, *Crematogaster* spp., *Hoplocampa minuta*, *Hoplocampa testudinea*, *Monomorium pharaonis*, *Solenopsis geminata*, *Solenopsis invicta*, *S. richteri*, *S. xyloni*, *Pogonomyrmex barbatus*, *Pogonomyrmex californicus*, *Dasymutilla occidentalis*,  
15 *Bombus* spp. *Vespula squamosa*, *Paravespula vulgaris*, *P. pennsylvanica*, *P. germanica*, *Dolichovespula maculata*, *Vespa crabro*, *Polistes rubiginosa*, *Camponotus floridanus*, and *Linepithium humile*,
- crickets, grasshoppers, locusts (*Orthoptera*), e.g. *Acheta domestica*, *Forficula auricularia*, *Gryllotalpa gryllotalpa*, *Locusta migratoria*, *Melanoplus bivittatus*, *Melanoplus femurrubrum*, *Melanoplus mexicanus*, *Melanoplus sanguinipes*, *Melanoplus spretus*, *Nomadacris septemfasciata*, *Schistocerca americana*, *Schistocerca peregrina*, *Stauronotus maroccanus* and *Tachycines asynamorus*,  
20
- fleas (*Siphonaptera*), e.g. *Ctenocephalidea felis*, *C. canis*, *Xenopsylla cheopis*, *Pulex irritans*, *Tunga penetrans*, and *Nosopsyllus fasciatus*  
25
- silverfish, firebrat (*Tysanura*), e.g. *Lepisma saccharina* and *Thermobia domestica*  
30
- lice (*Phthiraptera*), e.g. *Pediculus humanus capitis*, *Pediculus humanus corporis*, *Pythirus pubis*, *Haematopinus eurytetrus*, *Haematopinus suis*, *Linognathus vituli* and *Solenopotes capillatus*  
35
- Mesostigmata*, e.g. *Ornithonyssus bacoti* and *Dermanyssus gallinae*,  
*Prostigmata*, e.g. *Pymotes tritici*,  
*Astigmata*, e.g. *Acarus siro*,  
40
- ticks (*Ixodida*), e.g. *Phipicephalus sanguineus*,

## 11

*Saturniidae*, e.g. *Automeris io*,

Urticating caterpillars (*Lymantriidae*), e.g. *Lymantria dispar* and *Euproctis chrysorrhoea*,

5

*Pyralididae*, e.g. *Plodia interpunctella*, and *Ephestia kuhniella*,

*Gelechiidae*, e.g. *Sitotroga cerealella*,

10 *Meloidae*, e.g. *Epicauta fabricii* and *E. vittana*,

The compounds of formula I and compositions containing them are particularly useful for the control of pests of the orders Isoptera, Blattaria (Blattodea), Diptera, Dermaptera, Hemiptera, Thysanoptera, Hymenoptera, Orthoptera, Siphonaptera, Thysanura, and Phthiraptera.

15

Moreover, the compounds of formula I and compositions containing them are especially useful for the control of Isoptera, Diptera, Blattaria (Blattodea), Hymenoptera, and Siphonaptera.

20

In particular, the compounds of formula I and compositions containing them are useful for the control of Isoptera (*hodotermitidae*, *kalotermitidae*, *rhinotermitidae*, *termitidae*), Diptera (*tipulidae*, *blephariceridae*, *deuterophlebiidae*, *pachyneuridae*, *trichceridae*, *culicidae*, *simuliidae*, *ceratopogonidae*, *chironomidae*, *tabanidae*, *rhagionidae*, *mydidae*, *hilarimorphidae*, *tephritidae*, *braulidae*, *chloropidae*, *tethinidae*, *muscidae*, *caliphoridae*, *oestridae*, *sarcophagidae*, *rhinophoridae*, *tachnidae*, *hippoboscidae*), Blattaria/Blattodea (*cryptocercidae*, *blattidae*, *polyphagidae*, *blattellidae*), Hymenoptera (*xyelidae*, *argidae*, *cimbicidae*, *tenthredinidae*, *anaxyelidae*, *cephidae*, *aphidiidae*, *formicidae*, *vespoidae*, *sphécidae*), and Siphonaptera (*pulicidae*, *rhopalopsyllidae*, *ceratophyllidae*).

25

30

For use according to the present invention, the compounds I can be converted into the customary formulations, e.g. solutions, emulsions, suspensions, dusts, powders, pastes and granules. The use form depends on the particular purpose; it is intended to ensure in each case a fine and uniform distribution of the compound according to the invention.

35

The formulations are prepared in a known manner, e.g. by extending the active ingredient with solvents and/or carriers, if desired using emulsifiers and dispersants.

40

Solvents/auxiliaries, which are suitable, are essentially:

- water, aromatic solvents (for example Solvesso products, xylene), paraffins (for example mineral fractions), alcohols (for example methanol, butanol, pentanol, benzyl alcohol), ketones (for example cyclohexanone, gamma-butyrolactone),

## 12

pyrrolidones (NMP, NOP), acetates (glycol diacetate), glycols, fatty acid dimethylamides, fatty acids and fatty acid esters. In principle, solvent mixtures may also be used.

- 5       - carriers such as ground natural minerals (e.g. kaolins, clays, talc, chalk) and ground synthetic minerals (e.g. highly disperse silica, silicates); emulsifiers such as nonionic and anionic emulsifiers (e.g. polyoxyethylene fatty alcohol ethers, alkylsulfonates and arylsulfonates) and dispersants such as lignin-sulfite waste liquors and methylcellulose.
- 10      Suitable surfactants are alkali metal, alkaline earth metal and ammonium salts of lignosulfonic acid, naphthalenesulfonic acid, phenolsulfonic acid, dibutyl-naphthalenesulfonic acid, alkylarylsulfonates, alkyl sulfates, alkylsulfonates, fatty alcohol sulfates, fatty acids and sulfated fatty alcohol glycol ethers, furthermore condensates of sulfonated naphthalene and naphthalene derivatives with
- 15      formaldehyde, condensates of naphthalene or of naphthalenesulfonic acid with phenol and formaldehyde, polyoxyethylene octylphenyl ether, ethoxylated isooctylphenol, octylphenol, nonylphenol, alkylphenyl polyglycol ethers, tributylphenyl polyglycol ether, tristearylphenyl polyglycol ether, alkylaryl polyether alcohols, alcohol and fatty alcohol/ethylene oxide condensates, ethoxylated castor oil, polyoxyethylene alkyl
- 20      ethers, ethoxylated polyoxypropylene, lauryl alcohol polyglycol ether acetal, sorbitol esters, lignin-sulfite waste liquors and methylcellulose.

- Substances which are suitable for the preparation of directly sprayable solutions, emulsions, pastes or oil dispersions are mineral oil fractions of medium to high boiling
- 25      point, such as kerosene or diesel oil, furthermore coal tar oils and oils of vegetable or animal origin, aliphatic, cyclic and aromatic hydrocarbons, for example toluene, xylene, paraffin, tetrahydronaphthalene, alkylated naphthalenes or their derivatives, methanol, ethanol, propanol, butanol, cyclohexanol, cyclohexanone, isophorone, strongly polar solvents, for example dimethyl sulfoxide, N-methylpyrrolidone and water.

30

Powders, materials for spreading and dusts can be prepared by mixing or concomitantly grinding the active substances with a solid carrier.

- Granules, for example coated granules, impregnated granules and homogeneous
- 35      granules, can be prepared by binding the active ingredients to solid carriers. Examples of solid carriers are mineral earths such as silica gels, silicates, talc, kaolin, attaclay, limestone, lime, chalk, bole, loess, clay, dolomite, diatomaceous earth, calcium sulfate, magnesium sulfate, magnesium oxide, ground synthetic materials, fertilizers, such as, for example, ammonium sulfate, ammonium phosphate, ammonium nitrate, ureas, and
- 40      products of vegetable origin, such as cereal meal, tree bark meal, wood meal and nutshell meal, cellulose powders and other solid carriers.

## 13

In general, the formulations comprise from 0.01 to 95% by weight, preferably from 0.1 to 90% by weight, of the active ingredient. The active ingredients are employed in a purity of from 90% to 100%, preferably 95% to 100% (according to NMR spectrum).

5

The following are examples of formulations: 1. Products for dilution with water

A Soluble concentrates (SL)

10 10 parts by weight of a compound according to the invention are dissolved in water or in a water-soluble solvent. As an alternative, wetters or other auxiliaries are added. The active ingredient dissolves upon dilution with water.

B Dispersible concentrates (DC)

15 20 parts by weight of a compound according to the invention are dissolved in cyclohexanone with addition of a dispersant, for example polyvinylpyrrolidone. Dilution with water gives a dispersion.

C Emulsifiable concentrates (EC)

20 15 parts by weight of a compound according to the invention are dissolved in xylene with addition of calcium dodecylbenzenesulfonate and castor oil ethoxylate (in each case 5% strength). Dilution with water gives an emulsion.

D Emulsions (EW, EO)

25 40 parts by weight of a compound according to the invention are dissolved in xylene with addition of calcium dodecylbenzenesulfonate and castor oil ethoxylate (in each case 5% strength). This mixture is introduced into water by means of an emulsifier (Ultraturrax) and made into a homogeneous emulsion. Dilution with water gives an emulsion.

30 E Suspensions (SC, OD)

In an agitated ball mill, 20 parts by weight of a compound according to the invention are comminuted with addition of dispersant, wetters and water or an organic solvent to give a fine active ingredient suspension. Dilution with water gives a stable suspension of the active ingredient.

35

F Water-dispersible granules and water-soluble granules (WG, SG)

40 50 parts by weight of a compound according to the invention are ground finely with addition of dispersants and wetters and made into water-dispersible or water-soluble granules by means of technical appliances (for example extrusion, spray tower, fluidized bed). Dilution with water gives a stable dispersion or solution of the active ingredient.



## 14

G Water-dispersible powders and water-soluble powders (WP, SP)

75 parts by weight of a compound according to the invention are ground in a rotor-stator mill with addition of dispersant, wetters and silica gel. Dilution with water gives a stable dispersion or solution with the active ingredient.

5

2. Products to be applied undiluted

H Dustable powders (DP)

5 parts by weight of a compound according to the invention are ground finely and mixed intimately with 95% of finely divided kaolin. This gives a dustable product.

10

I Granules (GR, FG, GG, MG)

0.5 parts by weight of a compound according to the invention is ground finely and associated with 95.5% carriers. Current methods are extrusion, spray drying or the fluidized bed. This gives granules to be applied undiluted.

15

J ULV solutions (UL)

10 parts by weight of a compound according to the invention are dissolved in an organic solvent, for example xylene. This gives a product to be applied undiluted.

20

The active ingredients can be used as such, in the form of their formulations or the use forms prepared therefrom, eg. in the form of directly sprayable solutions, powders, suspensions or dispersions, emulsions, oil dispersions, pastes, dustable products, materials for spreading, or granules, by means of spraying, atomizing, dusting, spreading or pouring. The use forms depend entirely on the intended purposes; it is intended to ensure in each case the finest possible distribution of the active ingredients according to the invention.

25

Aqueous use forms can be prepared from emulsion concentrates, pastes or wettable powders (sprayable powders, oil dispersions) by adding water. To prepare emulsions, pastes or oil dispersions, the substances, as such or dissolved in an oil or solvent, can be homogenized in water by means of a wetter, tackifier, dispersant or emulsifier. Alternatively, it is possible to prepare concentrates composed of active substance, wetter, tackifier, dispersant or emulsifier and, if appropriate, solvent or oil, and such concentrates are suitable for dilution with water.

30  
35

The active ingredient concentrations in the ready-to-use products can be varied within relatively wide ranges. In general, they are from 0.0001 to 10%, preferably from 0.01 to 1%.

40

## 15

The active ingredients may also be used successfully in the ultra-low-volume process (ULV), it being possible to apply formulations comprising over 95% by weight of active ingredient, or even to apply the active ingredient without additives.

- 5 Various types of oils, wetters, adjuvants, herbicides, fungicides, other pesticides, or bactericides may be added to the active ingredients, if appropriate just immediately prior to use (tank mix). These agents usually are admixed with the agents according to the invention in a weight ratio of 1:10 to 10:1.

- 10 The compounds of formula I are effective through both contact (via soil, glass, wall, bed net, carpet, plant parts or animal parts), and ingestion (bait, or plant part).

According to a preferred embodiment of the invention, the compounds of formula I are prepared into a bait preparation.

15

For use against ants, termites, wasps, flies, mosquitos, crickets, or cockroaches, compounds of formula I are preferably used in a bait composition.

- 20 The bait can be a liquid, a solid or a semisolid preparation (e.g. a gel). Solid baits can be formed into various shapes and forms suitable to the respective application e.g. granules, blocks, sticks, disks. Liquid baits can be filled into various devices to ensure proper application, e.g. open containers, spray devices, droplet sources, or evaporation sources. Gels can be based on aqueous or oily matrices and can be formulated to particular necessities in terms of stickyness, moisture retention or aging characteristics.

25

- The bait employed in the composition is a product which is sufficiently attractive to incite insects such as ants, termites, wasps, flies, mosquitos, crickets etc. or cockroaches to eat it. The attractiveness can be manipulated by using feeding stimulants or sex pheromones. Food stimulants are chosen, for example, but not exclusively, from animal and/or plant proteins (meat-, fish- or blood meal, insect parts, egg yolk), from fats and oils of animal and/or plant origin, or mono-, oligo- or polyorganosaccharides, especially from sucrose, lactose, fructose, dextrose, glucose, starch, pectin or even molasses or honey. Fresh or decaying parts of fruits, crops, plants, animals, insects or specific parts thereof can also serve as a feeding stimulant. Sex pheromones are known to be more insect specific. Specific pheromones are described in the literature and are known to those skilled in the art.

30

35

- 40 Formulations of compounds of formula I as aerosols (e.g. in spray cans), oil sprays or pump sprays are highly suitable for the non-professional user for controlling pests such as flies, fleas, ticks, mosquitos or cockroaches. Aerosol recipes are preferably composed of the active compound, solvents such as lower alcohols (e.g. methanol, etha-

## 16

- nol, propanol, butanol), ketones (e.g. acetone, methyl ethyl ketone), paraffin hydrocarbons (e.g. kerosenes) having boiling ranges of approximately 50 to 250 °C, dimethylfomaamide, N-methylpyrrolidone, dimethyl sulphoxide, aromatic hydrocarbons such as toluene, xylene, water, furthermore auxiliaries such as emulsifiers such as sorbitol
- 5 monooleate, oleyl ethoxylate having 3-7 mol of ethylene oxide, fatty alcohol ethoxylate, perfume oils such as ethereal oils, esters of medium fatty acids with lower alcohols, aromatic carbonyl compounds, if appropriate stabilizers such as sodium benzoate, amphoteric surfactants, lower epoxides, triethyl orthoformate and, if required, propellants
- 10 nitrous oxide, or mixtures of these gases.

The oil spray formulations differ from the aerosol recipes in that no propellants are used.

- 15 The compounds of formula I and its respective compositions can also be used in mosquito and fumigating coils, smoke cartridges, vaporizer plates or long-term vaporizers and also in moth papers, moth pads or other heat-independent vaporizer systems.

- 20 The compounds of formula I are also suitable for the treatment of seeds. Conventional seed treatments include suspension concentrates FS and wettable powders WS.

The compounds of formula I can also be formulated into pour-on formulations.

- 25 The following list of pesticides together with which the compounds according to the invention can be used, is intended to illustrate the possible combinations, but not to impose any limitation:

- 30 Organophosphates: Acephate, Azinphos-methyl, Chlorpyrifos, Chlorfenvinphos, Diazinon, Dichlorvos, Dicrotophos, Dimethoate, Disulfoton, Ethion, Fenitrothion, Fenthion, Isoxathion, Malathion, Methamidophos, Methidathion, Methyl-Parathion, Mevinphos, Monocrotophos, Oxydemeton-methyl, Paraoxon, Parathion, Phenthoate, Phosalone, Phosmet, Phosphamidon, Phorate, Phoxim, Pirimiphos-methyl, Profenofos, Prothiofos, Sulprofos, Triazophos, Trichlorfon;

- 35 Carbamates: Alanycarb, Benfuracarb, Carbaryl, Carbosulfan, Fenoxycarb, Furathiocarb, Indoxacarb, Methiocarb, Methomyl, Oxamyl, Pirimicarb, Propoxur, Thiodicarb, Triazamate;

- 40 Pyrethroids: Bifenthrin, Cyfluthrin, Cypermethrin, Deltamethrin, Esfenvalerate, Ethofenprox, Fenpropathrin, Fenvalerate, Cyhalothrin, Lambda-Cyhalothrin, Permethrin, Silafluofen, Tau-Fluvalinate, Tefluthrin, Tralomethrin, Zeta-Cypermethrin;

## 17

Arthropod growth regulators: a) chitin synthesis inhibitors: benzoylureas: Chlorflua-  
zuron, Diflubenzuron, Flucycloxuron, Flufenoxuron, Hexaflumuron, Lufenuron, Novalu-  
ron, Teflubenzuron, Triflumuron; Buprofezin, Diofenolan, Hexythiazox, Etoxazole,  
Clofentazine; b) ecdysone antagonists: Halofenozide, Methoxyfenozide, Tebufenozide;  
5 c) juvenoids: Pyriproxyfen, Methoprene, Fenoxycarb; d) lipid biosynthesis inhibitors:  
Spirodiclofen;

Various: Abamectin, Acequinocyl, Amitraz, Azadirachtin, Bifenazate, Cartap, Chlor-  
fenapyr, Chlordimeform, Cyromazine, Diafenthiuron, Dinetofuran, Diofenolan, Ema-  
10 mectin, Endosulfan, Ethiprole, Fenazaquin, Fipronil, Formetanate, Formetanate hydro-  
chloride, Hydramethylnon, Imidacloprid, Indoxacarb, Metaflumizon, Pyridaben, Py-  
metrozine, Spinosad, Sulfur, Tebufenpyrad, Thiamethoxam, and Thiocyclam.

The insects may be controlled by contacting the target parasite/pest, its food supply,  
15 habitat, breeding ground or its locus with a pesticidally effective amount of compounds  
of or compositions of formula I.

"Locus" means a habitat, breeding ground, plant, seed, soil, area, material or environ-  
ment in which a pest or parasite is growing or may grow.  
20

In general, "pesticidally effective amount" means the amount of active ingredient  
needed to achieve an observable effect on growth, including the effects of necrosis,  
death, retardation, prevention, and removal, destruction, or otherwise diminishing the  
occurrence and activity of the target organism. The pesticidally effective amount can  
25 vary for the various compounds/compositions used in the invention. A pesticidally ef-  
fective amount of the compositions will also vary according to the prevailing conditions  
such as desired pesticidal effect and duration, weather, target species, locus, mode of  
application, and the like.

30 The compounds of formula I and its compositions can be used for protecting wooden  
materials such as trees, board fences, sleepers, etc. and buildings such as houses,  
outhouses, factories, but also construction materials, furniture, leathers, fibers, vinyl  
articles, electric wires and cables etc. from ants and/or termites, and for controlling ants  
and termites from doing harm to crops or human being (e.g. when the pests invade into  
35 houses and public facilities). The compounds of formula I are applied not only to the  
surrounding soil surface or into the under-floor soil in order to protect wooden materials  
but it can also be applied to lumbered articles such as surfaces of the under-floor con-  
crete, alcove posts, beams, plywoods, furniture, etc., wooden articles such as particle  
boards, half boards, etc. and vinyl articles such as coated electric wires, vinyl sheets,  
40 heat insulating material such as styrene foams, etc. In case of application against ants  
doing harm to crops or human beings, the ant controller of the present invention is ap-

## 18

plied to the crops or the surrounding soil, or is directly applied to the nest of ants or the like.

- 5 The compounds of the invention can also be applied preventively to places at which occurrence of the pests is expected.

- 10 The compounds of formula I may be also used to protect growing plants from attack or infestation by pests by contacting the plant with a pesticidally effective amount of compounds of formula I. As such, "contacting" includes both direct contact (applying the compounds/compositions directly on the pest and/or plant - typically to the foliage, stem or roots of the plant) and indirect contact (applying the compounds/compositions to the locus of the pest and/or plant).

- 15 In the case of soil treatment or of application to the pests dwelling place or nest, the quantity of active ingredient ranges from 0.0001 to 500 g per 100 m<sup>2</sup>, preferably from 0.001 to 20 g per 100 m<sup>2</sup>.

- 20 Customary application rates in the protection of materials are, for example, from 0.01 g to 1000 g of active compound per m<sup>2</sup> treated material, desirably from 0.1 g to 50 g per m<sup>2</sup>.

- 25 For use in bait compositions, the typical content of active ingredient is from 0.001 weight % to 15 weight %, desirably from 0.001 weight % to 5% weight % of active compound.

- For use in spray compositions, the content of active ingredient is from 0.001 to 80 weights %, preferably from 0.01 to 50 weight % and most preferably from 0.01 to 15 weight %.

- 30 For use in treating crop plants, the rate of application of the active ingredients of this invention may be in the range of 0.1 g to 4000 g per hectare, desirably from 25 g to 600 g per hectare, more desirably from 50 g to 500 g per hectare.

- 35 For use in treating seeds, the typical rate of application is from 1 g to 500 g per kilogram of seeds, desirably from 2 g to 300 g per kilogram of seeds, more desirably from 10 g to 200 g per kilogram of seeds.

- 40 This invention also provides a method for treating, controlling, preventing and protecting warm-blooded animals, including humans, and fish against infestation and infection by pests of the orders Siphonaptera, Hymenoptera, Hemiptera, Blattaria, Phthiraptera, and Diptera, which comprises orally, topically or parenterally administering or applying to said animals an pesticidally effective amount of compounds of formula I.

5 The invention also provides a process for the preparation of a composition for treating, controlling, preventing or protecting a warm-blooded animal or a fish against infestation or infection by pests of the Siphonaptera, Hymenoptera, Hemiptera, Blattaria, Phthiraptera, and Diptera orders which comprises a pesticidally effective amount of a compound having the formula I.

10 The above method is particularly useful for controlling and preventing infestations and infections in warm-blooded animals such as cattle, sheep, swine, camels, deer, horses, poultry, rabbits, goats, dogs and cats as well as humans.

15 Infestations in warm-blooded animals and fish including, but not limited to, lice, biting lice, ticks, nasal bots, keds, biting flies, muscoid flies, flies, myiasitic fly larvae, chiggers, gnats, mosquitoes and fleas may be controlled, prevented or eliminated by the compounds of formula I. The compounds of formula I are especially useful for combating pests of the following orders:

20 *Siphonaptera*, e.g. *Ctenocephalidea felis*, *C. canis*, *Xenopsylla cheopis*, *Pulex irritans*, *Tunga penetrans*, and *Nosopsyllus fasciatus*,

25 *Hymenoptera*: *Solenopsis* spp., *S. invicta*, *S. richteri*, *S. xyloni*, *S. geminata*, *Pogomyrmex* spp., *P. barbatus*, *P. californicus*, *Dasymutilla occidentalis*, Africanized honey bees, *Bombus* spp., *Vespula squamosa*, *Paravespula vulgari*, *P. pennsylvanica*, *P. germanica*, *Dolichovespula maculata*, *Vespa crabro*, and *Polistes rubiginosa*,

30 *Hemiptera*: *Cimex lectularius*, *C. hemipterus*, *Reduvius senilis*, *Triatoma* spp., and *Arilus critatus*,

*Blattaria*: *Blattella germanica*, *B. asahinae*, *Blatta orientalis*, *Periplaneta Americana*, *P. fuliginosa*, *P. brunnea*, and *P. australasiae*,

*Phthiraptera*: *Pediculus humanus corporis*, *P. h. capitis*, and *Pthirus pubis*,

35 *Diptera*: *Aedes aegypti*, *A. albopictus*, *Anopheles crucinas*, *An. albimanus*, *An. Gambiae*, *An. freeborni*, *An. leucosphyrus*, *An. minimus*, *An. quadrimaculatus*, *Culex nigripalpus*, *C. quinquefasciatus*, *C. tarsalis*, *Culiseta inornata*, *C. melanura*, *Mansonia titilanus*, *Psorophora columbiae*, *P. discolor*, *Prosimulium mixtum*, *Simulium vittatum*, *Chrysops discalis*, *C. silacea*, *C. atlanticus*, *Tabanus atratus*, *T. lineola*, *T. similes*, *Culicoides furens*, *Leptoconops torrens*, *Phlebotomus argentipes*, *Stomoxys calcitrans*,  
40 *Glossina* spp., *G. palpalis*, *G. fuscipes*, *G. tachinoides*, *Musca domestica*, *Sarcophaga haemorrhoidalis*, *Calliphora vicina*, *Hippelates* spp., *Dermatobia hominis*, *Chrysomya bezziana*, and *Cochliomyia hominivorax*.

## 20

For oral administration to warm-blooded animals, the formula I compounds may be formulated as animal feeds, animal feed premixes, animal feed concentrates, pills, solutions, pastes, suspensions, drenches, gels, tablets, boluses and capsules. In addition, the formula I compounds may be administered to the animals in their drinking water. For oral administration, the dosage form chosen should provide the animal with 0.01 mg/kg to 100 mg/kg of animal body weight per day of the formula I compound.

Alternatively, the formula I compounds may be administered to animals parenterally, for example, by intraruminal, intramuscular, intravenous or subcutaneous injection. The formula I compounds may be dispersed or dissolved in a physiologically acceptable carrier for subcutaneous injection. Alternatively, the formula I compounds may be formulated into an implant for subcutaneous administration. In addition the formula I compound may be transdermally administered to animals. For parenteral administration, the dosage form chosen should provide the animal with 0.01 mg/kg to 100 mg/kg of animal body weight per day of the formula I compound.

The formula I compounds may also be applied topically to the animals in the form of dips, dusts, powders, collars, medallions, sprays and pour-on formulations. For topical application, dips and sprays usually contain 0.5 ppm to 5,000 ppm and preferably 1 ppm to 3,000 ppm of the formula I compound. In addition, the formula I compounds may be formulated as ear tags for animals, particularly quadrupeds such as cattle and sheep.

The formula I compounds of this invention may also be used in combination or conjunction with one or more other parasitocidal compounds including, but not limited to, anthelmintics, such as benzimidazoles, piperazine, levamisole, pyrantel, praziquantel and the like; endectocides such as avermectins, milbemycins and the like; ectoparasitocides such as arylpyrroles, organophosphates, carbamates, gamma-butyric acid inhibitors including fipronil, pyrethroids, spinosads, imidacloprid and the like; insect growth regulators such as pyriproxyfen, cyromazine and the like; and chitin synthase inhibitors such as benzoylureas including flufenoxuron.

#### Examples of action against pests

#### 35 Test Methodology

1. Activity against argentine ant, harvester ant, acrobat ant, carpenter ant, fire ant, house fly, stable fly, flesh fly, yellowfever mosquito, house mosquito, malaria mosquito, German cockroach, cat flea, and brown dog tick via glass contact

Glass vials (20 ml scintillation vials) were treated with 0.5 ml of a solution of active ingredient in acetone. Each vial was rolled uncapped for ca. 10 minutes to allow the a.i.

## 21

to completely coat the vial and to allow for full drying of the acetone. Insects or ticks were placed into each vial. The vials were kept at 22 °C and were observed for treatment effects at various time intervals. Results are presented in Table 1.

- 5     2.     Activity against argentine ant, acrobat ant, carpenter ant, fire ant, and eastern subterranean termite via soil contact

10     For ants, tests were conducted in Petri dishes. A thin layer of 1 % agar in water was dispensed into the dishes and Florida sandy soil was spread over the agar (5 g for the small dishes and 11 g for the larger dishes). The active ingredient was dissolved in acetone and dispensed over the sand. Dishes were vented to evaporate the acetone, infested with ants, and covered. A 20% honey water solution was placed in each dish. The dishes were maintained at 22°C and observed for mortality at various time intervals.

15     For termites, a thin layer of 1% agar was dispensed into Petri dishes. A thin layer of pre-treated soil was spread over the agar. For soil treatment, the active ingredient was diluted in acetone on a weight-to-weight basis and incorporated into 100 g of soil. The soil was placed in a jar and vented for 48 hours. The moisture level of the soil was brought to field capacity by adding 7 ml of water. Termite workers were introduced into each dish. A small piece of filter paper was placed into each dish after 1 day as a food source, and additional water was added as needed to maintain soil moisture. Test dishes were held at a dark incubator at 25°C and 80% room humidity. Termites were observed daily for mortality (dead or unable to stand upright and showing only weak movement). Results are shown in Table 1.

- 30     3.     Activity against argentine ant, acrobat ant, carpenter ant, fire ant, house fly, eastern subterranean termite, formosan subterranean termite, and German cockroach via bait

35     For argentine ant, acrobat ant, and carpenter ant, tests were conducted in Petri dishes. Ants were given a water source, and then were starved of a food source for 24 hours. Baits were prepared with either 20% honey/water solutions or ground cat chow. Active ingredient in acetone was added to the bait. 0.2 ml of treated honey water solution or 150 mg of treated cat chow, placed in a cap, was added to each dish. The dishes were covered and maintained at a temperature of 22°C. The ants were observed for mortality daily. Results are shown in Table 1.

40     For the fire ants, corn grit was used as a bait matrix. Corn grit bait was prepared using a mixture of defatted corn grit (80%), soybean oil (19.9%), acetone, and the active ingredient (0.1%). Petri dishes were supplied with a water source. Fire ant adults were



## 22

placed into each dish. The next day, 250 mg of bait in bait containers was placed into the dishes. The ants were observed for mortality daily. Results are shown in Table 1.

For house flies. Bait tests were conducted with adults aged 2-5 days post-emergence.

- 5 Active ingredient in acetone was applied to a bait matrix consisting of a 1:1 mixture of powdered milk and sugar which was then allowed to dry. Assays were conducted in jars with 250 mg of bait in a pan placed in the bottom of each jar. House flies were placed into the bait jars which were covered. The test jars were held at 22°C. Test jars were observed at 4 and 24 hours after treatment for knockdown and 2 and 3 days after treatment for knockdown (death plus morbidity (unable to stay upright) plus intoxication (able to stand but unable to fly). Results are shown in Table 1.

- 15 For termites, active ingredient in acetone was applied to filter papers. % a.i. were calculated on basis of the weight of the filter paper. Acetone only was applied for untreated controls. Treated papers were vented to evaporate the acetone, moistened with ml water, and placed Petri dishes with sand. Water was added during the test as needed. Bioassays were conducted with one treated filter and ca. 30 termite workers per test dish. Test dishes were maintained at 25°C and 85% room humidity and observed daily for mortality (dead or moribund insects) or intoxication. Dead or moribund insects were removed daily. Results are shown in Table 1.

- 25 For cockroaches, plastic roach boxes with ventilated lids were used as test arenas. The top 3-4 cm of the arenas was treated with Vaseline and mineral oil to prevent roaches from escaping. Water was provided as needed. The bait was prepared using ground cat chow, and the active ingredient in acetone was incorporated on a weight- to-weight ratio. The treated chow was allowed to dry. The cockroaches were placed in the boxes and starved for 24 hours prior to bait introduction. 0.03 grams of bait per box were placed in a weigh boat. The boxes were maintained at 22°C and observed daily for mortality of the cockroaches. Results are shown in Table 1.

30

#### 4. Activity against yellowfever mosquito, southern house mosquito, and malaria mosquito larvae via water treatment

- 35 Well plates were used as test arenas. The active ingredient was dissolved in acetone and diluted with water to obtain the concentrations needed. The final Solutions containing appr. 1% acetone were placed into each well. Approximately 10 mosquito larvae (4<sup>th</sup>-instars) in 1 ml water were added to each well. Larvae were fed one drop of liver powder each day. The dishes were covered and maintained at 22°C. Mortality was recorded daily and dead larvae and live or dead pupae were removed daily. At the end of the test remaining live larvae were recorded and percent mortality was calculated. Results are shown in Table 1.

40

Each test was replicated at least 3 times.

23

## Results

Tests conducted with compounds of formula I-1 and I-2 showed the following results:

5

Table 1. Activity against various species.

Pest Common Name	Pest Latin Name	Rate	Days or Hours to achieve 100% mortality
Activity via glass contact			
argentine ant	<i>Linepithium humile</i>	10 ppm	1-2 days
harvester ant	<i>Pogonomyrmex californicus</i>	10 ppm	2-3 days
acrobat ant	<i>Crematogaster spp.</i>	10 ppm	1-2 days
carpenter ant	<i>Camponotus floridanus</i>	10 ppm	1 day
fire ant	<i>Solenopsis invicta</i>	10 ppm	4 hours
house fly	<i>Musca domestica</i>	10 ppm	4 hours
stable fly	<i>Stomoxys calcitrans</i>	10 ppm	4 hours
flesh fly	<i>Sarcophaga sp.</i>	10 ppm	4 hours
yellowfever mosquito	<i>Aedes aegypti</i>	10 ppm	4 hours
house mosquito	<i>Culex quinquefasciatus</i>	0.5 ppm	4 hours
malaria mosquito	<i>Anopheles albimanus</i>	1 ppm	1 day
german cockroach	<i>Blattella germanica</i>	100 ppm	5-24 hours
cat flea	<i>Ctenocephalides felis</i>	100 ppm	2 days
brown dog tick	<i>Rhipicephalus sanguineus</i>	10 ppm	3-5 days
Activity via soil contact			
argentine ant	<i>Linepithium humile</i>	0.1w%	1-2 days
acrobat ant	<i>Crematogaster spp.</i>	0.01w%	2 days
carpenter ant	<i>Camponotus floridanus</i>	0.01w%	1 day
fire ant	<i>Solenopsis invicta</i>	0.01w%	1 day
subterranean termite	<i>Reticulitermes flavipes</i>	0.005w%	1-3 days
Activity via bait			
argentine ant	<i>Linepithium humile</i>	1.0w%	2-3 days
acrobat ant	<i>Crematogaster spp.</i>	1.5w%	6-7 days
carpenter ant	<i>Camponotus floridanus</i>	2.0w%	1-3 days
fire ant	<i>Solenopsis invicta</i>	0.7w%	3 days
house fly	<i>Musca domestica</i>	0.1w%	3 hours
subterranean termite	<i>Reticulitermes flavipes</i>	0.1w%	1-2 days
formosan termite	<i>Coptotermes formosanus</i>	0.1w%	5 days


## 24

Pest Common Name	Pest Latin Name	Rate	Days or Hours to achieve 100% mortality
german cockroach	<i>Blattella germanica</i>	0.3w%	1-2 days
Activity via water treatment			
yellowfever mosquito	<i>Aedes aegypti</i>	10 ppm	2 days
house mosquito	<i>Culex quinquefasciatus</i>	10 ppm	1 day
malaria mosquito	<i>Anopheles albimanus</i>	1.0 ppm	1 day

25

Claims:


1. The use of compounds of formula I



5

wherein

Q is



10

$\text{X}^1$  is chlorine, bromine, or fluorine;

15

$\text{R}^1, \text{R}^2$  are each independently hydrogen,  $\text{C}_1\text{-C}_{10}$ -alkyl,  $\text{C}_3\text{-C}_{10}$ -alkenyl,  $\text{C}_3\text{-C}_{10}$ -alkynyl, or  $\text{C}_3\text{-C}_{12}$ -cycloalkyl,  $\text{C}_1\text{-C}_6$ -alkylamino, di( $\text{C}_1\text{-C}_6$ -alkyl)-amino,  $\text{C}_1\text{-C}_6$ -alkylcarbonylamino,  $\text{C}_1\text{-C}_6$ -alkylsulfonyl, or  $\text{C}_1\text{-C}_6$ -alkylsulfinyl, wherein the carbon atoms in these groups may be substituted with

20

1 to 3 halogen, hydroxy, nitro, cyano, amino, mercapto,  $\text{C}_1\text{-C}_6$ -alkoxy,  $\text{C}_1\text{-C}_6$ -haloalkoxy,  $\text{C}_1\text{-C}_6$ -alkylthio,  $\text{C}_1\text{-C}_6$ -haloalkylthio,  $\text{C}_1\text{-C}_6$ -alkylsulfonyl,  $\text{C}_1\text{-C}_6$ -alkylsulfinyl,  $\text{C}_1\text{-C}_6$ -haloalkylsulfonyl,  $\text{C}_1\text{-C}_6$ -haloalkylsulfinyl, or  $\text{C}_3\text{-C}_6$ -cycloalkyl which may be substituted with 1 to 3  $\text{R}^{\#}$  groups, or

25

$\text{R}^{\#}$  is halogen, cyano, nitro, hydroxy, mercapto, amino,  $\text{C}_1\text{-C}_6$ -alkoxy,  $\text{C}_2\text{-C}_6$ -alkenyloxy,  $\text{C}_2\text{-C}_6$ -alkynyloxy,  $\text{C}_1\text{-C}_6$ -haloalkoxy,  $\text{C}_1\text{-C}_6$ -alkylthio, or  $\text{C}_1\text{-C}_6$ -haloalkylthio,  $\text{C}_1\text{-C}_6$ -alkylsulfonyl,  $\text{C}_1\text{-C}_6$ -alkylsulfinyl,  $\text{C}_1\text{-C}_6$ -alkylamino, di( $\text{C}_1\text{-C}_6$ -alkyl)-amino,  $\text{C}_1\text{-C}_6$ -alkylcarbonyl,  $\text{C}_1\text{-C}_6$ -alkoxycarbonyl, or di( $\text{C}_1\text{-C}_6$ -alkyl)aminocarbonyl;

30


formyl,  $\text{C}_1\text{-C}_6$ -alkylcarbonyl,  $\text{C}(=\text{O})\text{NR}^{\text{a}}\text{R}^{\text{b}}$ ,  $\text{CO}_2\text{R}^{\text{c}}$ ,  $\text{R}^{\text{d}}$ ,  $\text{R}^{\text{e}}$ , phenyl which may be substituted with 1 to 3  $\text{R}^{\#}$  groups, or pyridyl which may be substituted with 1 to 3  $\text{R}^{\#}$  groups,

35

$\text{R}^{\text{a}}, \text{R}^{\text{b}}, \text{R}^{\text{c}}$  are each independently hydrogen or  $\text{C}_1\text{-C}_4$ -alkyl which may be substituted with 1 to 3 groups  $\text{R}^{\#}$ ;


$\text{R}^{\text{d}}$  is  $\text{NR}^{\text{i}}\text{R}^{\text{j}}$  or

26




5  $R^i, R^j$  are each independently hydrogen or  $C_1$ - $C_4$ -alkyl which may be substituted with 1 to 3 groups  $R''$ ;  
 $p, m$  are each independently 0, 1, 2, or 3, with the proviso that  $p$  and  $m$  are not both 0.  
 $X$  is oxygen, sulfur, amino,  $C_1$ - $C_4$ -alkylamino, or phenylamino, or, if  $p$  is 0 then  $X$  can also be phenoxy or  $C_1$ - $C_6$ -alkoxy;  
 $r$  is 0 or 1;

10  $R^a$  is



15  $R^k, R^a$  are each independently hydrogen or  $C_1$ - $C_4$ -alkyl which may be substituted with 1 to 3 groups  $R''$ ; or

20  $R^1$  and  $R^2$  may be taken together to form a ring represented by the structure



$p, m$  are 1, 2 or 3;

$X'$  is oxygen, sulfur, amino,  $C_1$ - $C_4$ -alkylamino, phenylamino, or methylene;

25  $Z$  is  $C_1$ - $C_4$ -alkyl or phenyl;

30  $R^3$  is hydrogen,  $C_1$ - $C_{10}$ -alkyl,  $C_2$ - $C_{10}$ -alkenyl,  $C_2$ - $C_{10}$ -alkynyl,  $C_3$ - $C_{12}$ -cycloalkyl, wherein the carbon atoms in these groups may be partially or fully halogenated or substituted with

1 to 3 cyano, nitro, hydroxy, mercapto, amino,  $C_1$ - $C_6$ -alkyl,  $C_3$ - $C_6$ -cycloalkyl,  $C_1$ - $C_6$ -alkoxy,  $C_1$ - $C_6$ -alkylamino, di( $C_1$ - $C_6$ -alkyl)-amino,  $C_1$ - $C_6$ -alkylthio,  $C_1$ - $C_6$ -alkylsulfonyl, or  $C_1$ - $C_6$ -alkylsulfinyl groups, wherein the carbon atoms in these groups may be substituted by

35 1 to 3 halogen atoms, a 5- to 6-membered aromatic ring system which may contain 1 to 4 heteroatoms selected from

## 27

- 5 oxygen, sulfur and nitrogen and which may be substituted with any combination of 1 to 5 halogen atoms, 1 to 3 C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-alkylthio, C<sub>1</sub>-C<sub>6</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, nitro, or cyano groups, wherein the carbon atoms in these groups may be substituted by 1 to 3 halogen atoms, or
- 10 phenoxy, which may be substituted with any combination of 1 to 5 halogen atoms, 1 to 3 C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-alkylthio, C<sub>1</sub>-C<sub>6</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, nitro, or cyano groups, wherein the carbon atoms in these groups may be substituted by 1 to 3 halogen atoms, or
- 15 a 3- to 6-membered saturated or partially unsaturated ring system which contains 1 to 3 heteroatoms selected from oxygen, sulfur and nitrogen and which may be substituted with any combination of 1 to 5 halogen atoms, 1 to 3 C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-alkylthio, C<sub>1</sub>-C<sub>6</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, nitro, or cyano groups, wherein the carbon atoms in these groups may be substituted by 1 to 3 halogen atoms,
- 20
- 25 a 3- to 6-membered saturated or partially unsaturated ring system which contains 1 to 3 heteroatoms selected from oxygen, sulfur and nitrogen and which is unsubstituted or substituted with any combination of 1 to 5 halogen atoms, 1 to 3 C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-alkylthio, C<sub>1</sub>-C<sub>6</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, C<sub>1</sub>-C<sub>6</sub>-haloalkoxy, nitro, or cyano groups, wherein the carbon atoms in these groups may be substituted by 1 to 3 halogen atoms;
- 30
- 35 R, R<sup>4</sup> are each independently hydrogen or C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkylaminocarbonyl, or di(C<sub>1</sub>-C<sub>6</sub>-alkyl)-aminocarbonyl, wherein the carbon atoms in the these groups may be substituted with 1 to 3 groups R<sup>#</sup>;
- A is C-R<sup>5</sup> or N;
- B is C-R<sup>6</sup> or N;
- W is C-R<sup>7</sup> or N;
- 40 with the proviso that one of A, B and W is other than N;

$R^5$ ,  $R^6$ ,  $R^7$  are each independently hydrogen, halogen, nitro, cyano, amino, mercapto, hydroxy,  $C_1$ - $C_{10}$ -alkyl,  $C_2$ - $C_{10}$ -alkenyl,  $C_2$ - $C_{10}$ -alkynyl,  $C_3$ - $C_6$ -cycloalkyl,  $C_1$ - $C_6$ -alkoxy,  $C_1$ - $C_6$ -alkylamino, di( $C_1$ - $C_6$ -alkyl)-amino,  $C_1$ - $C_6$ -alkylthio,  $C_1$ - $C_6$ -alkylsulfonyl, or  $C_1$ - $C_6$ -alkylsulfinyl, wherein the carbon atoms in these groups may be substituted with 1 to 3 groups  $R^{\#}$

a 5- to 6-membered aromatic ringsystem which may contain 1 to 4 heteroatoms selected from oxygen, sulfur and nitrogen and which may be substituted with any combination of 1 to 5 halogen atoms, 1 to 3  $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -haloalkyl,  $C_1$ - $C_6$ -alkylthio,  $C_1$ - $C_6$ -haloalkylthio,  $C_1$ - $C_6$ -alkylsulfonyl,  $C_1$ - $C_6$ -alkylsulfinyl,  $C_1$ - $C_6$ -haloalkylsulfonyl,  $C_1$ - $C_6$ -haloalkylsulfinyl,  $C_1$ - $C_6$ -alkoxy,  $C_1$ - $C_6$ -haloalkoxy, mercapto, hydroxy, amino, nitro, or cyano groups, wherein the carbon atoms in these groups may be substituted with 1 to 3 groups  $R^{\#}$ ;

$Y$  is hydrogen, halogen, cyano, nitro, amino, hydroxy, mercapto,  $C_1$ - $C_6$ -alkyl,  $C_2$ - $C_{10}$ -alkenyl,  $C_2$ - $C_{10}$ -alkynyl,  $C_3$ - $C_6$ -cycloalkyl,  $C_1$ - $C_6$ -alkoxy,  $C_1$ - $C_6$ -alkylamino, di( $C_1$ - $C_6$ )-alkylamino,  $C_1$ - $C_6$ -alkylthio,  $C_1$ - $C_6$ -alkylsulfonyl, or  $C_1$ - $C_6$ -alkylsulfinyl, wherein the carbon atoms in these groups may be substituted with 1 to 3 groups  $R^{\#}$ ;

$n$  is 0, 1, or 2;

or the enantiomers or diastereomers, salts or esters thereof,

for combating non-crop pests.



2. The use according to claim 1 of compounds of formula I as defined in claim 1 wherein

A is  $C-R^5$ ;

B is  $C-R^6$ ; and

W is  $C-R^7$ .

3. The use according to claim 1 wherein the compound of formula I is a compound of formula I-1 or I-2.



I-2


4. The use according to any one of claims 1 to 3 wherein the non-crop pests are selected from the Isoptera, Diptera, Blattaria (Blattodea), Hymenoptera, and Siphonaptera orders.
- 5 5. The use of compounds as defined in any one of claims 1 to 3 for protecting wooden materials against pests from the Isoptera and Hymenoptera orders.
6. A method for controlling non-crop pests comprising contacting the pests or their food supply, habitat or breeding grounds with a pesticidally effective amount of a compound of formula I as defined in any one of claims 1 to 3.
- 10 7. A bait composition which comprises a compound of formula I as defined in any one of claims 1 to 3 and a food stimulant.
- 15 8. A method for treating, controlling, preventing or protecting a warm-blooded animal or a fish against infestation or infection by pests of the orders *Siphonaptera*, *Hymenoptera*, *Hemiptera*, *Blattaria*, *Phthiraptera* and *Diptera* which comprises orally, topically or parenterally administering or applying to said animal or fish a pesticidally effective amount of a compound having the formula I as defined in any one of claims 1 to 3.
- 20 9. A process for the preparation of a composition for treating, controlling, preventing or protecting a warm-blooded animal or a fish against infestation or infection by the *Siphonaptera*, *Hymenoptera*, *Hemiptera*, *Blattaria*, *Phthiraptera* and *Diptera* orders which comprises a pesticidally effective amount of a compound of formula I as defined in any one of claims 1 to 3.
- 25




30

## Abstract

The use of compounds of formula I



wherein Q is




5

 $X^1$  is halogen;

$R^1$  and  $R^2$  are each H, or optionally substituted alkyl, alkenyl, alkynyl, cycloalkyl, alkylamino, dialkyl-amino, alkylcarbonylamino, alkylsulfonyl, or alkylsulfinyl, or  $R^1$  and  $R^2$  may be taken together to form a ring represented by the structure below, wherein p,m

are 1, 2 or 3,  $X^1$  is O, S, NH, alkylamino, phenylamino, or methylene, Z is alkyl or phenyl;

10

 $R^3$  is H or optionally substituted alkyl, alkenyl, alkynyl, cycloalkyl,

$R$ ,  $R^4$  are each H, or optionally substituted alkyl, alkoxy, carbonyl, alkylaminocarbonyl, or dialkylaminocarbonyl,

15

A is  $\text{C-R}^5$  or N; B is  $\text{C-R}^6$  or N; W is  $\text{C-R}^7$  or N; with the proviso that one of A, B and W must be other than N;

$R^5$ ,  $R^6$ ,  $R^7$  are each H, halogen,  $\text{NO}_2$ , CN,  $\text{NH}_2$ , SH, OH, or optionally substituted alkyl, alkenyl, alkynyl, cycloalkyl, alkoxy, alkylamino, dialkylamino, alkylthio, alkylsulfonyl, or alkylsulfinyl, or a 5- to 6-membered optionally substituted aromatic ring system which may contain 1 to 4 heteroatoms selected from oxygen, sulfur and nitrogen

20

Y is H, halogen, CN,  $\text{NO}_2$ ,  $\text{NH}_2$ , OH, SH, or optionally substituted alkyl, alkenyl, alkynyl, cycloalkyl, alkoxy, alkylamino, dialkylamino, alkylthio, alkylsulfonyl, or alkylsulfinyl, n is 0, 1, or 2;

25

or the enantiomers or diastereomers, salts or esters thereof, for combating pests and compositions comprising compounds I.